

Robert T. Sataloff, Associate Editor

Reentry Following COVID-19: Concerns for Singers

David Meyer, John Nix, Lynn Holding, Allen Henderson, Tom Carroll, Jeremy Faust, Christine Petersen



David Meyer



John Nix



Lynn Holding



Allen Henderson



Tom Carroll



Jeremy Faust



Christine Petersen

INTRODUCTION

THE TARGET OF THIS PAPER IS CAPTURED in its subtitle: “Concerns for Singers.” While the COVID-19 pandemic has wreaked devastation upon disparate types of people around the world, singers, and those who teach and collaborate with them, are a vulnerable cohort within this larger global health crisis. This vulnerability is due to a number of facts about the SARS-CoV-2 virus, beginning with its three main routes of transmission as outlined by the U.S. Centers for Disease Control and Prevention (CDC): (1) inhalation of very fine respiratory droplets and aerosol particles; (2) the deposition of virus-containing droplets and particles on exposed mucous membranes in the mouth, nose, or eye by direct splashes and sprays; and (3) touching mucous membranes with hands that have been soiled either directly by virus-containing respiratory fluids or indirectly by touching surfaces with virus on them.¹

The CDC has further explained that “the risk of SARS-CoV-2 infection varies according to the amount of virus to which a person is exposed” and notes that the two main variables concerning this amount are *distance* and *time*.² Regarding the former, how far away an individual is from the respiratory droplets of an infected person, combined with the observation that the concentration of the virus is diluted by both gravity (heavier drops fall) and mixture with air, illustrates why the CDC states that “the available evidence continues to demonstrate that existing recommendations to prevent SARS-CoV-2 transmission remain effective. These include physical distancing.”³ The parameter of time exerts a similar dilution of viral viability and potency.

Yet using those same parameters of distance and time, the CDC also notes that “transmission of SARS-CoV-2 from inhalation of virus in the air farther than six feet from an infectious source *can* occur [emphasis added] under certain preventable circumstances,” one of which is “Prolonged exposure to these conditions, typically more than 15 minutes.”⁴ The CDC report goes on to note two other factors that increase the risk of SARS-CoV-2 infection.

- Enclosed spaces with inadequate ventilation or air handling within which the concentration of exhaled respiratory fluids, especially very fine droplets and aerosol particles, can build-up in the air space.

Journal of Singing, published ahead of print
<https://doi.org/10.53830/VAPD6085>
 Copyright © 2021
 National Association of Teachers of Singing

- Increased exhalation of respiratory fluids if the infectious person is engaged in physical exertion or increases the intensity of their voice (e.g., exercising, shouting, singing).⁵

It has been shown that there are significantly higher emission rates for singing compared to mouth breathing and speaking.⁶ This finding plus the CDC report indicates that the typical voice teacher who instructs singers on a back to back hourly schedule in a one on one setting, often in a small room with poor ventilation, may be especially vulnerable to COVID infection, particularly from unvaccinated students and collaborators (e.g., pianists). Indeed, voice teachers in these situations could be considered “sitting ducks” for multiple respiratory illnesses, from the common cold to influenza, but the virulence of the SARS-CoV-2 virus and the mortality rate of COVID-19 illness are without modern precedent: in the United States, a country with 331,449,281 persons⁷, 33,558,862 cases of COVID-19 and 602,275 fatalities⁸ have been reported as of June 22, 2021. These figures indicate that **one in ten** persons contracted the virus in the US, and of those who contracted COVID-19, **one in 56** died from the illness.

This ongoing global health crisis has added a heightened level of urgency to a list of concerns for singers. As noted by this team of authors in a previous article with the same subtitle, that list of concerns includes the COVID-19 *sequelae* most likely to disrupt or permanently damage those functions that are necessary for singing.⁹

The degree to which voice teachers can protect themselves from communicable diseases like COVID-19 begins with the amount of personal agency they wield. Self-employed singing teachers possess a large degree of autonomy regarding their work environment, including the degree and length of time to which they may be exposed to the SARS-CoV-2 virus; so do singers who volunteer for a community or church choir. Notwithstanding the emotionally difficult choices involved with avoidance or complete cessation of singing activities, the fact remains that such people do have choices. Yet teaching faculty at music conservatories or teachers in commercial music companies may have few, if any, choices about their teaching environments.

In anticipation of a return to in-person teaching and learning, a growing number of colleges and universities are requiring vaccination against COVID-19 for faculty,

staff, and students who intend to pursue in-person studies on campus this fall.¹⁰ Yet reentry to live interaction for institutionally based teachers and singers is less clear cut than this requirement might suggest, first by allowing vaccine hesitant students to decline the vaccine, ostensibly for health or religious reasons. A number of other factors complicate reentry, including vaccine hesitancy among nonstudent staff and faculty colleagues, changing CDC guidelines and recommendations, foreign students who have had vaccines not authorized by the FDA, and university administrations and commercial studios so anxious to return to in-person teaching and learning to recoup the economic losses of the past year that they have issued directives to employees forbidding them from inquiring about vaccination status among their students and collaborators.

This article aims to provide resources to help singers safely and ethically navigate a return to in-person instruction by considering the following:

- (1) Vaccination and Voice Teachers
 - Monitoring Local Levels of SARS-CoV2 Incidence
 - Safety of Vaccination
 - Vaccination and SARS-CoV-2 Variants
- (2) Mask Use Recommendations
- (3) Aerosol Risk Update
- (4) Environmental Risk Factors and Mitigation
 - Risk Factors in Studio Voice Instruction
 - Environmental Risk Mitigation
 - Increasing Air Changes Per Hour
 - CO₂ Meters
 - Air Changes Between Lessons/Rehearsals and Lesson Duration
- (5) FERPA, HIPAA, and Vaccine Hesitancy
- (6) Psychological Risks Associated with the Pandemic and Reentry
 - The General Psychological Impact of the COVID-19 Pandemic
 - The Physiological Impact of Stress, Anxiety, and Trauma on Voice
 - Resources and Advice in Support of Mental Health
- (7) Risk Assessment

VACCINATION AND VOICE TEACHERS

[Disclaimer: the following vaccination information and risk estimates are for Pfizer-BioNTech and Moderna

complete (2 shot) vaccinations. Johnson and Johnson Janssen, AstraZeneca, Sputnik V, and Sinovac vaccines will have differing levels of efficacy against emerging variants. For more information, consult your healthcare provider.]

A high rate of vaccination is the only path toward the safest possible “new normal” within 2021. We can never achieve zero risk for a viral infection that is likely to remain in low levels of circulation for the foreseeable future. However, vaccination can and does render SARS-CoV-2 comparable to many other respiratory infections that we routinely encounter in its likelihood to cause illness or death. Without vaccination, SARS-CoV-2 is without a doubt a far deadlier and more dangerous virus than those other pathogens, such as influenza or respiratory syncytial virus.¹¹ Safe in-person teaching and learning is best facilitated by high rates of SARS-CoV-2 vaccination in any given community, as this decreases viral spread and the potential creation of variant viruses and lowers the frequency of severe disease.¹²

Voice teachers are at higher risk for *all* viruses that spread through droplets or are in the air because of small studio size and the resultant concentration of exposure to droplets or virus aerosols in that studio.¹³ However, because most teach one person at a time, the mathematical odds of exposure for solo voice teachers in any given day, week, or month will remain low (as compared to other higher-population density locations such as public transit or large packed lecture halls). While a small number of vaccinated people can become infected from solo voice lesson settings, an even smaller number is likely to go on to experience substantial disease.¹⁴

Monitoring Local Levels of SARS-CoV-2 Incidence

There are multiple user friendly resources to monitor the local SARS-CoV-2 incidence (number of infections over a given period of time) and get a sense of one’s personal risk of exposure, usually per 100,000 people. A popular source has been the COVID-19 Dashboard provided by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (available at <https://coronavirus.jhu.edu/map.html>). The *New York Times* has provided similar maps and data about both case incidence and vaccination rates, using the Johns Hopkins data as well as locally sourced data (available at <https://www.nytimes.com/interactive/2021/world/>

[covid-cases.html](#)). In addition, many states and counties within the United States have data dashboards, but these are beginning to be removed in some locations; users are advised to check whether the information has been recently updated.

Safety of Vaccination

The technology used in all of the currently approved vaccines in the United States is based on decades of peer reviewed, validated science, including use of mRNA vaccines.¹⁵ The evidence provided from these studies in animals, clinical trials in people for previous vaccines, and the data from hundreds of millions of people now vaccinated is that the risk of an adverse event, or a “severe vaccine reaction” like anaphylaxis or death is less than 0.002%.¹⁶ The annual risk of death in an automobile crash or from a firearm related incident far exceeds this value.¹⁷ All of the published data to date indicate that COVID-19 vaccines approved for use in the United States are safe and effective during pregnancy.¹⁸ The impact of a COVID-19 infection on a pregnant woman, on the other hand, is notable. COVID-19 has been associated with a 25 to 30-fold increase in maternal mortality.¹⁹ Vaccination safely prevents infection and disease in pregnant women. In addition, some antibodies generated by vaccination are found in breast milk, which likely provides some degree of passive, short term protection to breastfed infants.²⁰

Vaccination and SARS-CoV-2 Variants

Some SARS-CoV-2 variants of concern may reduce the success rate of vaccinations; this reduction in success has not been confirmed for the Pfizer and Moderna vaccinations.²¹ However, if overall hospitalization rates remain low in a community, the composite risk among vaccinated persons is manifestly not different than historical risks we accept from other endemic respiratory viruses, most of which typically cause mild or even moderate self-limited disease in most targets, but which in some instances can cause serious disease (i.e., in those with other health risks). Given that, the opportunity for arts organizations to “return to normal” and “stay open” should not hinge on the presence of a small number of breakthrough cases, even serious ones, but rather on hospitalization patterns. If rates of hospitalization are at historically normal levels and vaccination rates are

high, live performing arts should remain safe. However, if variants emerge that evade the protection provided by vaccines such that there are once again spikes in hospitalization (in any age group), a return to stricter mitigation measures may become necessary.

MASK USE RECOMMENDATIONS

Persons who meet *all* the following criteria do not need to routinely wear masks in indoor group settings: those who are fully vaccinated, who do not have special risks (such as a profound immunocompromised state seen in those who are bone marrow transplant or solid organ transplant recipients who take antirejection medications),²² and who are living in areas with low COVID-19 incidence, and when the overall majority of the group is greater than 70% vaccinated.²³ While it is possible for vaccinated people to become infected, the likelihood of spread *from* such persons is low,²⁴ especially in asymptomatic infections (unlike unvaccinated asymptomatic infection, in which spread has been frequently documented).²⁵ Even though vaccinated individuals with breakthrough infections could theoretically spread the virus to a small number of people, vaccination of any potential contact provides a great deal of protection.²⁶

People who are vaccinated, but who have prolonged and frequent exposure to unvaccinated persons, may reasonably choose to wear a mask to prevent breakthrough infection to others. Mask wearing in this setting would be less necessary if unvaccinated persons are required to be routinely tested, with mandatory isolation or quarantine of any people testing positive as well as their contacts and/or in areas of very low incidence of COVID-19.²⁷ Rapid antigen tests are highly effective in detecting the presence of contagious virus in a test sample and provide information that is available in time to be safely relied upon for decision-making and removal of potentially infectious persons from the group.²⁸ The CDC and WHO recommend that unvaccinated people continue to wear masks and physically distance themselves when inside with others. Most recently, the WHO has recommended the continuation of mask-wearing by all individuals.²⁹ As the global public health body, this advice is grounded in the need to prevent spread of new viral “mutants” or variants of concern that can arise in locations with very low vaccination rates globally that

may be able to avoid or “escape” the protection provided by immune responses produced by vaccination.³⁰

This masking requirement, while most appropriate globally, should be considered necessary indoors in areas of the United States with local/county level COVID-19 vaccination rates of less than 70% of all adults, but is also strongly recommended in crowded locations elsewhere inside and out. It is best practice to have an informal conversation with students and colleagues about vaccination to establish what precautions should be taken to protect all parties. In addition, based on the remarkable downward trends of influenza globally as well as within the United States in the last year due to mask wearing and social distancing,³¹ healthcare professionals and public officials are indicating that to prevent having the flu or other serious viral infections (like RSV in small children), they are seriously considering regular mask use starting in the late fall and continuing through the high transmission period of February.³² Those who sing or are around people singing regularly might also take this into consideration as a wise precaution.

AEROSOL RISK UPDATE

As opposed to direct transmission through a handshake or other direct person to person contact, indirect transmission, and specifically airborne transmission, is now accepted as the dominant form of SARS-CoV-2 transmission.³³ The airborne route occurs during phonation, coughing, sneezing, and certain medical procedures, among other activities. There are three size ranges of particles that contribute to infection transmission: droplets, the largest particles, and by definition greater than 5 microns (μ); medium sized particles of 1 to 5 μ ; and small particles of 1 μ or smaller.³⁴ Airborne transmission can occur by any of these three particle sizes, however the medium sized particles pose the greatest risk, despite the recommended six feet of physical distancing being followed.³⁵ Room size, the number of infected people in a space, the activity being performed in the room, and room ventilation all play a role in risk of transmission and will be discussed elsewhere in this article.

Large droplets tend to dominate in a cough, as opposed to typical speech. Small particles, less than one μ , carry little virus but can float around for hours. Thus, medium particles contribute to the majority of disease transmis-

sion.³⁶ Less than five microns in size, the medium particles then dry and shrink in the air, concentrating their viral load as so-called “droplet nuclei” and float for hours in a space, thus creating the need for proper mitigation by those considering using that space.³⁷

Breathing, speaking, and singing lead to aerosolization of typically small particles that originate in the bronchioles and alveoli of the lungs. A fluid film burst occurs as we exhale, leading to these small particles being released from the lungs.³⁸ There is also some evidence that the larynx, the vocal folds, and possibly the oral cavity produce a majority of medium-sized particles through a similar method of fluid film burst, which is why speaking and singing contribute to the viral load in a closed space.³⁹ When speech or singing is louder, more aerosolization occurs.⁴⁰ Lung volume seems to directly correlate with expelled viral load: adolescents and adults have been shown to produce more aerosols than children.⁴¹

Face masks have been integral in mitigating the spread of the SARS-CoV-2 virus to some extent; however, there is no confirmatory conclusion of this due to conflicting data on the efficiency of wearing face masks to *prevent* the spread.⁴² We do know that strict social distancing and wearing masks reduces transmission and thus mitigates risk. Short range airborne transmission, meaning transmission through the air from one person to another, of droplets less than 100 μ likely is more common than typical droplet transmission when distances are 0.2 m or less during normal talking or 0.5 m or less during coughing.⁴³ Droplets will fall to the floor short of these distances if they are 100 μ or larger, whereas droplets and particles smaller than this size will move through the air to potentially infect another person in proximity.⁴⁴ Thus, six feet of distance may not be enough in certain situations.⁴⁵

It has been demonstrated that unmasked transmission can occur farther than six feet apart, up to 10 meters when the infected person is unmasked.⁴⁶ As previously discussed, there is some conflicting data on mask use, as it is also demonstrated that two meters of distancing is probably safe if both parties are masked.⁴⁷ Small and medium particles can move up to five meters, obviously longer than two meters or six feet of distance. As a caveat to this, Issakov et al. demonstrated that

there is a decrease in concentration of particles in the direction of airflow.⁴⁸

ENVIRONMENTAL RISK FACTORS AND MITIGATION

Risk Factors in Studio Voice Instruction

Studio voice instruction consists of one to one sessions between a singing teacher and a student, with the possible addition of a collaborative pianist in the teaching space. While this intimate setting allows artists to explore their potential without public scrutiny, this same intimacy can leave teachers, students, and other collaborators vulnerable to respiratory infections.⁴⁹ Singing, and in particular more robust types of singing that feature high sound pressure levels, has been shown to be a high aerosol generating activity.⁵⁰ Instruction occurs in various sized venues and ventilation conditions; teachers outside of academia may work in shared studios or in places of residence, while those in academic settings may teach in practice rooms (graduate assistants, adjunct faculty) or faculty offices. In the U.S., no standards exist regarding academic instruction room size; the major accreditation handbook only states “Space allotted to any music unit function must be adequate for the effective conduct of that function,” and “All instructional facilities shall be accessible, safe, and secure, and shall meet the standards of local fire and health codes.”⁵¹ As such, teachers, students, and pianists may interact in rooms too small for physical distancing measures and with variable or unknown ventilation. Both shared rooms and individual offices have a number of frequently touched surfaces, such as doors, piano keys, music stands, etc., which may serve as fomites. Teachers and pianists usually interact with multiple students per day, often in close succession, with each student having a unique health history and hygiene habits that are often not disclosed to the teacher or pianist.

Environmental Risk Mitigation

Given the previously mentioned conditions, teachers of singing, voice students, and pianists could be considered at greater risk for transmitting the SARS-CoV-2 virus and other upper respiratory contagious illnesses than the general population. However, several relatively low cost changes to the teaching and rehearsing environment

may reduce much of this increased risk. The changes include increasing the number of air changes per hour (ACH) in the teaching or rehearsal space; increasing the amount of outside fresh air brought directly into the room; using HEPA filtration units; using CO₂ meters to track room ventilation in real time; allowing air change time between lessons and rehearsals, and modifying lesson durations according to the vaccination status of the individuals involved and the community infection level. Each of these changes/interventions is detailed below.

Increasing Air Changes Per Hour. Air changes per hour (ACH) is defined as the volume of air added to or removed from a space in one hour divided by the volume of the space.⁵² Higher ACH values provide better dilution or removal of potential infectious aerosolized particles, reducing exposure time and therefore lowering risk.⁵³ Means for increasing air changes include natural ventilation with outside air, recirculated air that is filtered, and the use of portable high efficiency particle air (HEPA) filters.⁵⁴ Of these, admitting outside air through the opening of windows or outdoor air dampers is the most desirable,⁵⁵ although building design, external temperatures, and local air quality levels may preclude this. Modern buildings typically feature heating, ventilation, and air conditioning (HVAC) systems that introduce fresh air and exhaust a percentage of the recirculated air continuously. In the United States, recirculated air is filtered for particulate removal with no less than a MERV 8 filter.⁵⁶ Thus, the stated ACH in public buildings is a mix of fresh air changes and filtered air, with the filtered air considered as equivalent air changes based on the efficiency of the building's filtration.⁵⁷ Portable HEPA filtration units also are a viable, cost effective (\$100–250 as of July 1, 2021) and flexible means for increasing the total ACH. Filters meeting the HEPA standard must remove at least 99.97% of particles from the air that passes through with a diameter = 0.3 microns, with filtration efficiency increasing for particle diameters smaller and larger than 0.3 microns.⁵⁸ HEPA filters thus are theoretically capable of capturing small SARS-CoV-2 viral particles (0.1 microns) as well as medium sized particles and droplet nuclei (<5 microns).⁵⁹ A key means to quickly determine the suitability of a HEPA unit for a teaching space is the Clean Air Delivery Rate (CADR). By comparing the HEPA filter's CADR in cubic feet per

minute with the room's volume, an equivalent ACH value for the filter can be determined; this can be added to the fresh and recirculated ACH values to determine the total ACH for the space.

CO₂ Meters. Carbon dioxide (CO₂) meters can be used to measure carbon dioxide levels in parts per million in a teaching space. While CO₂ meters are not regarded as providing accurate information about the viral load in a room, they can provide cost effective (\$100–165 as of July 1, 2021) real-time information on the fresh air ventilation in a room, which can be an analog of the ACH rate and by extension the ability to dilute or remove potentially infectious aerosolized particles.⁶⁰ CO₂ levels rise with poor fresh air ventilation, an increased number of persons in a room, and increased physical activity level of persons in the room.⁶¹ Based on available evidence, the authors recommend a CO₂ level of 800 ppm be used as a threshold level of good air quality. When levels exceed 800 ppm, more fresh air should be admitted.⁶²

Air Changes Between Lessons/Rehearsals and Lesson Duration. A final means to make inexpensive environmental changes to reduce infection risk is allowing air changes between lessons/rehearsals⁶³ and reducing lesson/rehearsal durations.⁶⁴ The former provides time for the room ventilation to dilute or remove potential infectious particles, and the latter reduces the total number of particles emitted in a lesson. Teachers and pianists are advised to know the ACH of their teaching/rehearsal spaces. At 6 ACH, one air change occurs every 10 minutes. The authors recommend at least one air change (fresh air or filtered equivalent) after each lesson and advise limiting lesson lengths to 30 minutes if the singer is unvaccinated.⁶⁵

FERPA, HIPAA, AND VACCINE HESITANCY

There are two United States federal laws currently in the forefront of considerations among educators, administrators, studio owners, and others as they prepare for the lifting of COVID-19 restrictions and in-person collaboration. The *Health Insurance Portability and Accountability Act* of 1996 (HIPAA) was created to protect sensitive patient health information from being disclosed without the patient's consent or knowledge. The US Department of Health and Human Services

issued the companion *HIPAA Privacy Rule* to actually implement the requirements of HIPAA, which addresses several issues, including the use and disclosure of individuals' protected health information by entities subject to the Privacy Rule.

A major goal of the Privacy Rule is to ensure that individuals' health information is properly protected while allowing the flow of health information needed to provide and promote high quality health care and to protect the public's health and well-being. The Privacy Rule strikes a balance that permits important uses of information while protecting the privacy of people who seek care and healing.⁶⁶

The second federal law, the *Family Educational Rights and Privacy Act* (FERPA) protects the privacy of student education records and applies to all schools that receive funds from the US Department of Education.⁶⁷ The intersection of these two federal privacy laws is causing confusion for teachers and administrators in the era of COVID-19. It is rather simple to clarify the first of these (HIPAA) for voice teachers. Because HIPAA only applies to specific health care entities such as insurance providers, health care clearinghouses, health care providers and their business associates, voice teachers who do not work in a clinical environment on a medical team should not be concerned with violating HIPAA. If employers, supervisors, or even clients themselves attempt to use HIPAA as a justification to prevent a voice teacher from discussing vaccine hesitancy with voice students, they are in error. The only exception to this is voice teachers who work in a clinical environment.

FERPA is another misunderstood statute in the COVID-19 era. In March 2020, the US Department of Education issued a "Frequently Asked Questions (FAQs)" document to clarify issues about FERPA and COVID-19.⁶⁸ First, FERPA only applies to educational records maintained by an institution. Learning about a student's vaccination status through a social media post or casual conversation is not a violation of FERPA. Further, because the US Department of Health and Human Services declared COVID-19 a Public Health Emergency in January 2020, there are emergency provisions under the FERPA "health or safety emergency exception" that allow an educational agency to make determinations on a case by case basis inclusive of the

totality of circumstances. Teaching in one on one settings in studios is such a unique setting that a case could be made to share information.

Under the FERPA health or safety emergency exception, an educational agency or institution is responsible for making a determination, on a case-by-case basis, whether to disclose PII [personally identifiable information] from education records, and it may take into account the totality of the circumstances pertaining to the threat. . . . If the educational agency or institution determines that there is an articulable and significant threat to the health or safety of the student or another individual and that certain parties need the PII from education records, to protect the health or safety of the student or another individual, it may disclose that information to such parties without consent.⁶⁹

Some entities and institutions are issuing policies and guidelines claiming that asking persons (including students) about their vaccination status is a violation of one or both of these US privacy laws; some are even doing this without explicitly using HIPAA or FERPA as justifications, but vaguely referring to potential violations of privacy via "coercive or pressuring behavior."⁷⁰

At best, such policies may be in error, and at worst, are justifications used to wield control over important information and those who have access to it. Communications like these when sent to employees seem to indicate that there are no exceptions, while the reality is that there may be a number of allowable exceptions to privacy laws like FERPA. Indeed, the FERPA "health and safety emergency" exception is the mechanism within which justifiable exceptions are made.

Nevertheless, faculty should exercise sensitivity and diplomacy when inquiring about vaccine status among students, especially in institutional settings. Primarily these are concerns centered on religious and disability discrimination rather than protected health information. Therefore, it is important that singing teachers in each of the following settings fully understand the exceptions provided for under the FERPA "health or safety emergency exception."

Independent studio teachers are not subject to either HIPAA or FERPA, but should establish studio policies based on their setting and personal risk profile. They should know the current age limits for vaccination eli-

gibility and adjust their studio policies accordingly via effective communication with clients, and/or underage students and their parents or guardians, and monitor and enforce their policies when necessary.

Voice teachers and collaborative pianists at institutions without a vaccination requirement should continue promoting vaccination using advocacy messaging provided by their institution or professional associations. The joint statement issued by the NATS Voice Science Advisory Committee, the American Choral Directors Association, Chorus America, the Barbershop Harmony Society, the Performing Arts Medicine Association, the Pan American Vocology Association, Opera America, and the National Collegiate Choral Organization is one example.⁷¹ Almost all institutions have developed elaborate vaccine advocacy campaigns vetted by their legal offices and distributed using print and social media. Therefore, a faculty member may be viewed as an excellent team player by using these resources to advocate that everyone, including their private students, be vaccinated. Faculty should further note that there are multiple ways to discover the status of students while steering clear of privacy concerns; for example, faculty may negotiate vaccine status through an intermediary party, such as their campus student or employee health service who can provide an “all clear” rather than divulge individual’s health information.

Finally, it is likely that many teachers already know the status of most of their students either through students volunteering that information in conversation or on social media channels.

Unvaccinated or immunocompromised teachers and collaborative pianists at institutions should contact their Human Resources or appropriate office on campus to negotiate accommodation. It could be possible for such teachers to use the FERPA health or safety emergency exception to justify certain parameters of requested accommodations, such as: knowing the vaccination status of all students they are expected to teach; receiving the institution’s assurance of the safest possible work environment in terms of room size, ventilation, masking, lesson length, and/or CO2 monitoring or other factors impacting the safety of the teacher; and establishing protocols for each category of student (vaccinated, unvaccinated, and/or immunocompromised).

Voice teachers and pianists at institutions with a vaccination requirement may view FERPA with minimal concern because the institution is requiring all to be vaccinated. Many of these institutions are allowing exemptions for unvaccinated individuals based on health concerns or religious convictions. In such cases, teachers should work cooperatively to establish protocols for safe teaching in these unique settings as listed below for “Vaccinated teachers at institutions.”

Vaccinated teachers and pianists at institutions should realize they are significantly protected by being vaccinated. Nevertheless, due to their unique one on one teaching setting, often in small spaces, teachers in this setting are strongly encouraged to:

- follow the recommendations for “Voice teachers and collaborative pianists at institutions without a vaccination requirement” (above);
- request informed discussions with appropriate authorities on campus;
- prepare for such discussions by studying the FERPA “health or safety emergency exception” (previously referenced in this section);
- start the discussion with the claim that in order to provide a safe environment in these unique spaces, it is vitally important to know the vaccination status of each student they are expected to teach;
- provide a customized solution for each category of student (vaccinated, unvaccinated, and/or immunocompromised);
- mention the FERPA “health or safety emergency exception” as justification for your request of the release of PII (personally identifiable information).

These steps should open the door to discussion to protect the health and safety of all who enter the voice studio.

PSYCHOLOGICAL RISKS ASSOCIATED WITH THE PANDEMIC AND REENTRY

COVID-19 has posed serious direct threats to physical health and general wellbeing, and also has triggered psychological problems for many, including anxiety, depression, and post-traumatic stress disorder. While voice teachers are not (typically) licensed mental health-care providers, we must consider the psychological scars

of the COVID-19 epidemic as we return to in-person instruction. This section will examine the following: the general psychological impact of the COVID-19 pandemic, the physiological impact of stress, anxiety, and trauma, and resources and advice to support singers' mental health.

The General Psychological Impact of the COVID-19 Pandemic

Studies examining the psychological impact of the COVID-19 are emerging. Qiu et al. developed the COVID-19 Peritraumatic Distress Index (CPDI) and used it with a sample of over 53,000 persons in China.⁷² They examined the frequency of anxiety, depression, specific phobias, loss of social functioning, and other psychological factors related to the pandemic. Nearly a third of the study's respondents reported symptoms of peritraumatic distress during the pandemic lockdown in China. Similar levels of distress were reported in Italian,⁷³ French,⁷⁴ and Filipino studies,⁷⁵ Iranian,⁷⁶ and Brazilian⁷⁷ studies reported moderate to severe distress in over 60% of their participants. While CPDI levels in many of these studies declined when lockdown periods were lifted, meta-analyses of trauma literature have predictively linked severe levels of peritraumatic distress to subsequent psychopathology.⁷⁸

COVID-19 peritraumatic distress was more severe in certain populations. Studies found that persons 10–30 years of age experienced very high levels of distress,⁷⁹ possibly due to their increased consumption of social media⁸⁰ and news related to the outbreak.⁸¹ Gender also appears to have been a factor, with female identifying respondents showing significantly higher CPDI scores than male identifying counterparts.⁸² Previous studies also suggest that female identifying persons may be more likely to develop posttraumatic stress disorder and may be more generally vulnerable to the effects of stress.⁸³

Highly educated persons tended to have more severe COVID-19 peritraumatic distress.⁸⁴ American university students experienced high levels of depression (43.3%), high anxiety scores (45.4%), and high levels of PTSD symptoms (31.8%) in a study by Liu et al.⁸⁵ They reported that increased levels of loneliness, COVID-19-specific worry, and low distress tolerance were associated with clinical levels of depression, anxiety, and PTSD symptoms.⁸⁶ In a study of university students in China,

Tang et al. found that extreme pandemic fear was a significant risk factor for peritraumatic distress, followed by short sleep durations.⁸⁷ Alqudah et al. found that university students in Jordan experienced severe anxiety (40.6%) and mild to moderate anxiety (23.5%) due to the shift to distance learning, quarantine isolation, and increased student workload.⁸⁸

In short, the COVID-19 pandemic's psychological impact is unprecedented in the modern era; it is global in scope, has received extraordinary documentation due to smartphones and social media, and has created high levels of anticipatory anxiety due to the expected course and spread of the virus.⁸⁹ Horesh and Brown suggest it should therefore be treated as a new type of mass trauma, and hypothesize that it has exacerbated existing mental health disorders and created a new wave of stress-related disorders.⁹⁰ Persons who are particularly vulnerable include those who are young, identifying as female, university students, and those who frequently consume social media/news concerning COVID-19.⁹¹ This describes the majority of our collegiate singers.

The Physiological Impact of Stress, Anxiety, and Trauma on Voice

The vocal instrument is biological and physiological, and singers' voices are sensitive to the effects of stress. Psychologically, the voice is central to many singers' sense of self,⁹² and many find it difficult to decouple "who I am" from "what I do." This point was clearly described by Stohrer.

Singers are both the instrument and the player. Unable to put the instrument back in its case and leave it in the corner, a singer must carry it around, exposing it to . . . all the current viruses.⁹⁴

When identity and singing are inseparably linked, the loss of performance opportunities can cause considerable distress.

Stress has physical consequences. Prior to the pandemic it was estimated that 50% to 70% of all visits to a physician's office were due to stress related illness.⁹⁴ Stress can alter oral and vocal fold secretions, heart rate, and gastric acid production, and is associated with head and neck tension, decreased ability to concentrate, and chronic fatigue.⁹⁵ Vocal fold pathologies (e.g., reflux laryngitis and arytenoid irritation) can have etiologies

linked to increased gastric acid secretions caused by stress,⁹⁶ and high levels of anxiety may increase the risk of benign voice disorders.⁹⁷ Chronic stress has been linked to more dangerous problems such as asthma, depression of the immune system, and myocardial infarction.⁹⁸ Making matters worse, the risk of atherosclerotic cardiovascular disease is exacerbated by quarantine conditions due to physical inactivity.⁹⁹ Thus, the stressful conditions singers have experienced during the COVID-19 pandemic may have serious long lasting physical aftereffects.

Stress and anxiety can negatively affect control of voice and respiration.¹⁰⁰ Muscular spasms in the larynx have been observed due to the laryngeal nerve's sensitivity to emotional distress.¹⁰¹ Traumatic experiences (e.g., a global pandemic, the death of a family member, a health/financial crisis) can lead to voice symptoms without underlying physical pathology.¹⁰² Such symptoms were observed in a study of 313 Israeli collegiate professors during the COVID-19 pandemic. The authors found that psychological stress during the transition to online teaching was associated with elevated levels of voice problems. This was especially the case for those who reported high levels of psychological stress.¹⁰³ Monti et al. noted that emotional and psychological elements can have a profound effect on a singer's voice, and they encourage singing teachers to be mindful that technical difficulties may have deeply rooted emotional etiologies.¹⁰⁴

Trauma can also have a profound neurophysiological effect on singers. When the brain perceives a threat to survival (e.g., COVID-19 peritraumatic distress), a "fight/flight/freeze" response may engage that shuts down Broca's area in the left frontal lobe of the cortex.¹⁰⁵ This profoundly inhibits one's ability to put thoughts and feelings into words, and can even mimic the effects of a stroke or other physical lesion.¹⁰⁶ According to polyvagal theory, perceived threats to survival may also produce "a neural dissolution from systems of positive social behavior and social communication,"¹⁰⁷ making it difficult to feel connected to and safe around others.¹⁰⁸ Cues of safety and social engagement may ameliorate these neurophysiological symptoms of trauma.¹⁰⁹

Singers around the world have experienced trauma and loss related to the COVID-19 pandemic. This trauma may be manifested in social phobias, music

performance anxiety, depression, and grief. The authors of this paper hypothesize that epidemic levels of generalized anxiety, music performance anxiety, and depression may be seen in singers during the forthcoming re-entry phase of the pandemic.

Resources and Advice in Support of Mental Health

There are numerous resources now available to support the mental health of singers and teachers of singing. Music performance anxiety (MPA) has been studied extensively in the literature.¹¹⁰ In *The Musician's Mind: Teaching, Learning, and Performance in the Age of Brain Science*, Lynn Holding presents a useful review of the MPA literature and discusses standard psychotherapies, alternative therapies, and experimental therapies to ameliorate its symptoms.¹¹¹

Depression is a common medical condition to consider when working with singers. Nearly one in five people will suffer from major depression at some point in their lives,¹¹² and this prevalence may increase following the COVID-19 pandemic. Common speaking voice symptoms of depression include a flat affect, slowed rate of speech, decreased length of utterance, lengthy pauses, decreased pitch variability, mono-loudness, and frequent use of vocal fry.¹¹³ Discussing depression and other mental health issues can be awkward,¹¹⁴ but singers and teachers of singing are best served when these discussions are normalized. Pedagogues are encouraged to cultivate professional relationships with mental health care providers. As always, "when in doubt, refer out."

Integrating an awareness of and sensitivity to trauma in pedagogic practice may benefit singers as they return to in-person instruction. While trauma awareness initiatives are becoming more common in schools,¹¹⁵ similar initiatives are generally lacking in higher education.¹¹⁶ Literature on the therapeutic effects of singing, particularly with survivors of abuse and trauma, is an excellent starting place.¹¹⁷ Jess Baldwin's "Incorporating Basic Trauma Awareness into the Voice Lesson" also offers readers useful advice.¹¹⁸ She recommends teachers prioritize self-regulation (see also Van Eekelen et al.),¹¹⁹ develop a plan and communicate it, establish connection rituals, choose connection over productivity, validate students' feelings, break tasks into smaller segments, show appreciation for effort, focus on positive resources,

and be aware of the student's self-regulation (see also "student self-efficacy" in Crocco and Meyer)¹²⁰ while encouraging calm, fun, and connection in the lesson. Additional information on goal setting, journaling, and the use of psychology to build connection in the voice studio may be found in Meyer and Helling's "Voice Pedagogy: Practical Science in the Studio: 'No-Tech' Strategies."¹²¹

We all likely have experienced pandemic-induced loss. Many continue to suffer the lingering effects of trauma, anxiety, depression, and grief. Singing instructors must consider the psychological scars of the COVID-19 epidemic as we return to in-person instruction, **because we do not teach voice, we teach people**. When in doubt, refer students and colleagues to mental and medical healthcare providers, and consider building trauma awareness into standard pedagogic practice.

RISK ASSESSMENT

In a 2020 publication written by many of the present authors,¹²² several risk assessment tools and a decision assistance tool were provided. Here we provide an updated risk/decision guidance chart (see Table 1 below), which lists a number of moderately priced environmental and behavioral interventions available to the singing instruction community. Two levels of risk are shown, and references are provided for each intervention option. Other resources associated with a June 23, 2021 National Association of Teachers of Singing webinar led by the authors, "Reentry after COVID—Concerns for singers," can be found at https://www.nats.org/cgi/page.cgi/_article.html/Featured_Stories/_NATS_COVID_Resources_Page#webinars.

CONCLUSION

The SARS-CoV-2 pandemic has forced the singing voice community to undergo rapid and substantial change. Responses to the potentially life threatening situation have included several key elements not discussed in this article, chief among them being an explosion of remote learning technologies, a reassessment of instructional delivery methods (including synchronous in person and online teaching as well as asynchronous work), and new or expanded platforms for sharing live performances, all of which were undertaken to reduce the risks associ-

ated with close personal contact. As the U.S. and other Western countries continue to expand vaccination and slowly emerge from COVID-19 restrictions, we have reason for cautious optimism.¹²³ However, as they prepare to return to in-person instruction and performing, teachers, singers, and their collaborators should reexamine the unique risks that they collectively face at this point in the pandemic.

We close with a plea for full vaccination and continued patience (and, as warranted by local conditions, vigilance) by the singing voice community. For those teachers anxious about a return to in-person instruction, we urge actively controlling what can be controlled: first and foremost, one's own vaccination status; next, ensure that the risk level within the teaching environment or rehearsal studio is as low as possible; additionally, take proactive behavioral steps such as those widely in use prior to the availability of the vaccines, such as masking and distancing; finally, open a healthy and productive dialogue with students and colleagues about vaccination and other COVID-19 health issues, including mental/emotional ones. For those teachers who are eager for "business as usual," we suggest that a prudent approach is still warranted. The SARS-CoV-2 virus continues to mutate, and infections remain prevalent among the unvaccinated. There will likely be periods of greater and lesser public health restrictions in the months and years ahead in response to local conditions. We suggest maintaining the many hygiene habits acquired during the last 18 months, especially among the unvaccinated, not only for COVID-19 mitigation, but also for other upper respiratory illnesses that frequently affect the singing voice community. Simple behaviors can make a marked difference in reducing risks to teachers, singers, and collaborative pianists: vaccinate, ventilate, mitigate, and communicate.

ACKNOWLEDGEMENTS

The authors wish to thank the following professionals for their expert feedback during the creation of this article:

- Rebecca Gayle Morrison, Ph.D., Clinical Psychologist
- Sarah Van Orman MD, Associate Vice Provost for Student Health and Chief Health Officer, University of Southern California Student Health, and Clinical Professor of Family Medicine, Keck School of Medicine, USC

- Maura Copeland, Associate Vice President of Legal Affairs, Georgia Southern University
- Lesley Persily, LPC (retired)
- Ronald C. Scherer, Ph.D., Distinguished Research Professor, Department of Communication Sciences and Disorders, Bowling Green State University
- Dr. Robert Sataloff, Professor and Chairman, Department of Otolaryngology—Head and Neck Surgery, Senior Associate Dean for Clinical Academic Specialties, Drexel University College of Medicine

TABLE 1. Environmental and Behavioral Risk Assessment.

ENVIRONMENT	At Risk	Safer	Notes	References
Air changes per hour (ACH)	<3	6 or more	More air changes per hour removes aerosols from the teaching/rehearsal space, reducing exposure time not only to potential COVID infections, but also cold and flu viruses.	doi:10.1001/jama.2021.5053
Ability to open windows in teaching space	None	Yes	The ability to bring outside air directly into the teaching space is a key means to reduce the number of possibly infectious aerosol particles in the room.	https://www.cdc.gov/coronavirus/2019-ncov/community/ventilation.html ; https://osf.io/7rczy
HEPA filtration	None	Yes (with a room-size appropriate unit—sufficient CADR for volume of the room)	HEPA filtration can be a cost effective means to increase the number of equivalent air changes per hour in a teaching space that does not have windows.	JAMA and CDC references above; see also https://www.ashrae.org/file%20library/technical%20resources/covid-19/in-room-air-cleaner-guidance-for-reducing-covid-19-in-air-in-your-space-or-room.pdf ; https://pubmed.ncbi.nlm.nih.gov/33940414/ ; https://pubmed.ncbi.nlm.nih.gov/32662746/ ; https://pubmed.ncbi.nlm.nih.gov/32660218/ ; https://www.epa.gov/coronavirus/air-cleaners-hvac-filters-and-coronavirus-covid-19 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7711180/ ; https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8084223/
Room size	100 square feet or less (10' × 10')	200 square feet or more (14' × 14')	The ability to distance oneself from another person who is singing depends upon room size. Distancing can reduce exposure risks from directly expelled particles (larger droplets) containing COVID, cold, and flu viruses.	See a calculator tool at: https://indoor-covid-safety.herokuapp.com/ ; additional calculator available at https://www.mpic.de/4747361/risk-calculator?en ; https://www.jvoice.org/article/S0892-1997(20)30245-9/fulltext

(table continues)

TABLE 1. Environmental and Behavioral Risk Assessment (continued).

ENVIRONMENT	At Risk	Safer	Notes	References
CO₂ meter for real-time monitoring ventilation	None	CO ₂ meter in singing/rehearsal studio	CO ₂ meters provide a useful real-time analog to the number of local ACH. They do not provide any information about viral load, but they are a useful measure of the ventilation system's removal of expired air.	https://www.medrxiv.org/content/10.1101/2020.10.26.20218354v1.full); https://www.cdc.gov/coronavirus/2019-ncov/community/ventilation.html ; https://pubmed.ncbi.nlm.nih.gov/32028176/ ; https://www.health.state.mn.us/communities/environment/air/toxins/co2.html ; https://www.epa.gov/sites/production/files/2014-08/documents/indoorair20-247.pdf ; https://pubs.acs.org/doi/10.1021/acs.estlett.1c00183 ; https://www.dhs.wisconsin.gov/chemical/carbondioxide.htm
BEHAVIORAL STEPS	At Risk	Safer	Notes	References
Vaccination	Teacher, student, and accompanist are not vaccinated	Teacher, accompanists, and students are all fully vaccinated	All three FDA-authorized vaccines are readily available in the US. Vaccination is safe, effective, and free. Annual flu and pneumonia vaccines are also recommended.	https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/pfizer-biontech-covid-19-vaccine ; https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/moderna-covid-19-vaccine ; https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/janssen-covid-19-vaccine
Entrance screening	No symptom check for all individuals in the teaching/rehearsal space	Teachers, accompanists, and students all self-screen for COVID-19 symptoms before gathering	Screening apps include: https://covid19.apple.com/screening/ ; https://www.cdc.gov/screening/index.html ; https://www.webmd.com/coronavirus/coronavirus-assessment/default.htm .	https://www.doh.wa.gov/Portals/1/Documents/1600/coronavirus/Employervisitorscreeningguidance.pdf ; https://www.cdc.gov/coronavirus/2019-ncov/community/guidance-business-response.html#anchor_1609683211941
Mask usage	None	Yes for all unvaccinated persons	CDC recommendations include masking for all unvaccinated persons when gathering indoors. Vaccinated persons with reduced immune function may be encouraged to continue masking by their physicians. Masks can also reduce the risk of cold and flu virus transmission.	https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/participate-in-activities.html

(table continues)

TABLE 1. Environmental and Behavioral Risk Assessment (*continued*).

BEHAVIORAL STEPS	At Risk	Safer	Notes	References
Distancing	Guidelines not being followed	6 or more feet for unvaccinated persons	CDC distancing recommendations for vaccinated persons have been removed as of May 16, 2021; for unvaccinated persons, 6 feet or more is still recommended. Distancing can also help reduce the risk of cold and flu transmission.	https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/participate-in-activities.html
Lesson length	greater than 30 minutes	30 minutes maximum with unvaccinated persons	Recommendation based on research at the University of Colorado and the University of Maryland.	https://scholar.colorado.edu/concern/file_sets/9s161736t
Air change time between lessons	None	At least one air change	Increase if an unvaccinated person sings unmasked or if infection rates in community are rising.	doi:10.1001/jama.2021.5053
Aerosol producing activities (lip trills, raspberries, straw in water, vigorous fricatives, etc)	Yes, especially if unmasked or unvaccinated	Not used in lessons, but can be encouraged for solo practice in well-ventilated spaces.	Some SOVTs produce large amounts of droplets; masks may prevent the spread of droplets but not finer aerosols. Some 'singer friendly masks' may permit SOVT use in highly ventilated areas with vaccinated singers only. Vaccination reduces the chance of the singer shedding viral particles. Cold and flu viruses may also be spread through lip trill and raspberry use in group settings.	pending report from Colorado State University Bioaerosol Emission study – See https://smt.d.colostate.edu/reducing-bioaerosol-emissions-and-exposures-in-the-performing-arts/
Cleaning common surfaces between lessons	None	Yes, using alcohol or other anti-viral and anti-bacterial cleansers	Reduces the risk of exposure through fomites on hands.	https://www.who.int/westernpacific/emergencies/covid-19/information/transmission-protective-measures ; https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public
Teacher and students cleaning hands before lessons	None	Yes, using hot soapy water or alcohol-based cleansers	Reduces the risk of exposure through touching common surfaces and then touching the nose, eyes, and mouth.	https://www.who.int/westernpacific/emergencies/covid-19/information/transmission-protective-measures ; https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public
Age of students taught	Unvaccinated adults and children ineligible for vaccination	Vaccinated adolescents and adults	Adults produce more aerosols, but are more likely to be vaccinated; adolescents and children (especially under age 12) are less likely to be vaccinated.	https://journals.plos.org/plosone/article/authors?id=10.1371/journal.pone.0246819

NOTES

1. US Centers for Disease Control and Prevention (CDC), "Scientific Brief: SARS-CoV-2 Transmission, Updated May 7, 2021"; <https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/sars-cov-2-transmission.html> (accessed June 28, 2021).
2. Ibid.
3. Ibid.
4. Ibid.
5. Ibid.
6. Dirk Murbe, Martin Kriegel, Julia Lange, Hansjorg Rotheudt, and Mario Fleischer, "Aerosol emission is increased in professional singing" (August 26, 2020): 1; https://depositonce.tu-berlin.de/bitstream/11303/11491/5/muerbe_et_al_2020_aerosols-singing.pdf (accessed July 7, 2021).
7. US Census Bureau, "Table A. Apportionment Population, Resident Population, and Overseas Population: 2020 Census and 2010 Census"; <https://www2.census.gov/programs-surveys/decennial/2020/data/apportionment/apportionment-2020-tableA.pdf> (accessed July 7, 2021).
8. Johns Hopkins University, "COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU)"; <https://coronavirus.jhu.edu/map.html> (accessed June 22, 2021).
9. Lynn Holding, Thomas L Carroll, John Nix, Michael M. Johns, Wendy D. LeBorgne, and David Meyer, "COVID-19 After Effects: Concerns for Singers," *Journal of Voice* (in press; available online August 6, 2020): 1; <https://doi.org/10.1016/j.jvoice.2020.07.032> (accessed July 7, 2021).
10. Andy Thomason and Brian O'Leary, "Here's a List of Colleges That Will Require Students or Employees to be Vaccinated Against Covid-19," *Chronicle of Higher Education* online version (July 6, 2021); <https://www.chronicle.com/blogs/live-coronavirus-updates/heres-a-list-of-colleges-that-will-require-students-to-be-vaccinated-against-covid-19> (accessed July 7, 2021).
11. Lindsey R. Baden, Hana M. El Sahly, Brandon Essink, Karen Kotloff, Sharon Frey, Rick Novak, David Diemert, Stephen A. Spector, Nadine Rouphael, C. Buddy Creech, John McGettigan, Shishir Khetan, et al., for the COVE Study Group, "Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine," *New England Journal of Medicine* 384, no. 5 (February 4, 2021): 403–416; <https://doi.org/10.1056/NEJMoa2035389> (accessed July 7, 2021); Fernando P. Polack, Stephen J. Thomas, Nicholas Kitchin, Judith Absalon, Alejandra Gurtman, Stephen Lockhart, John L. Perez, Gonzalo Pérez Marc, Edson D. Moreira, Cristiano Zerbini, Ruth Bailey, Kena A. Swanson, et al., for the C4591001 Clinical Trial Group, "Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine," *New England Journal of Medicine* 383, no. 27 (December 31, 2020): 2603–2615; <https://doi.org/10.1056/NEJMoa2034577> (accessed July 7, 2021); Sarah Geoghegan, Anabella Erviti, Mauricio T. Caballero, Fernando Vallone, Stella M. Zanone, Juan Ves Losada, Alejandra Bianchi, Patricio L. Acosta, Laura B. Talarico, Adrian Ferretti, Luciano Alva Grimaldi, et al., "Mortality due to Respiratory Syncytial Virus. Burden and Risk Factors," *American Journal of Respiratory and Critical Care Medicine* 195, no. 1 (January 1, 2017): 96–103; <https://doi.org/10.1164/rccm.201603-0658OC> (accessed July 7, 2021); Jeremy S. Faust and Carlos del Rio, "Assessment of Deaths From COVID-19 and From Seasonal Influenza," *JAMA Internal Medicine* 180, no. 8 (May 14, 2020): 1045; <https://doi.org/10.1001/jamainternmed.2020.2306> (accessed July 7, 2021).
12. Noa Dagan, Noam Barda, Eldad Kepten, Oren Miron, Shay Perchik, Mark A. Katz, Miguel A. Hernán, Marc Lipsitch, Ben Reis, and Ran D. Balicer, "BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting," *New England Journal of Medicine* 384, no. 15 (April 15, 2021): 1412–1423; <https://doi.org/10.1056/NEJMoa2101765> (accessed July 7, 2021).
13. Zahra Noorimotlagh, Neemat Jaafarzadeh, Susana Silva Martínez, and Seyyed Abbas Mirzaee, "A systematic review of possible airborne transmission of the COVID-19 virus (SARS-CoV-2) in the indoor air environment," *Environmental Research* 193:110612 (February 2021); [doi:10.1016/j.envres.2020.110612](https://doi.org/10.1016/j.envres.2020.110612) (accessed July 7, 2021).
14. Dagan, Barda, Kepten, et al.
15. Thomas Schlake, Andreas Thess, Mariola Fotin-Mleczek, and Karl-Josef Kallen, "Developing mRNA-vaccine technologies," *RNA Biology* 9, no. 11 (October 12, 2012): 1319–1330; <https://doi.org/10.4161/rna.22269> (accessed July 7, 2021); Ivana Knezevic, Margaret A. Liu, Keith Peden, Tiequn Zhou, and Hye-Na Kang, "Development of mRNA Vaccines: Scientific and Regulatory Issues," *Vaccines* 9, no. 2 (January 23, 2021): 81; <https://doi.org/10.3390/vaccines9020081> (accessed July 7, 2021).
16. US Centers for Disease Control and Prevention, "Selected Adverse Events Reported after COVID-19 Vaccination, updated June 30, 2021"; <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/safety/adverse-events.html> (accessed July 9, 2021).
17. Insurance Information Institute, "Facts + statistics: mortality risk"; <https://www.iii.org/fact-statistic/facts-statistics-mortality-risk> (accessed July 9, 2021).

18. Tom T. Shimabukuro, Shin Y. Kim, Tanya R. Myers, Pedro L. Moro, Titilope Oduyebo, Lakshmi Panagiotakopoulos, Paige L. Marquez, Christine K. Olson, Ruiling Liu, Karen T. Chang, Sascha R. Ellington, Veronica K. Burkel, et al., for the CDC v-safe COVID-19 Pregnancy Registry Team, "Preliminary Findings of mRNA Covid-19 Vaccine Safety in Pregnant Persons," *New England Journal of Medicine* 384, no. 24 (April 21, 2021):2273–2282; <http://doi.org/10.1056/NEJMoa2104983> (accessed July 9, 2021).
19. Jose Villar, Shabina Ariff, Robert B. Gunier, et al., "Maternal and Neonatal Morbidity and Mortality Among Pregnant Women With and Without COVID-19 Infection: The INTERCOVID Multinational Cohort Study," *JAMA Pediatrics* Published online April 22, 2021; <https://doi.org/10.1001/jamapediatrics.2021.1050> (accessed July 9, 2021).
20. Sivan H. Perl, Atara Uzan-Yulzari, Hodaya Klainer, Liron Asiskovich, Michal Youngster, Ehud Rinott, and Ilan Youngster, "SARS-CoV-2–Specific Antibodies in Breast Milk After COVID-19 Vaccination of Breastfeeding Women," *Journal of the American Medical Association* 325, no. 19 (April 12, 2021):2013–2014; <https://doi.org/10.1001/jama.2021.5782> (accessed July 9, 2021).
21. Philip R. Krause, Thomas R. Fleming, Ira M. Longini, Richard Peto, Sylvie Briand, David L. Heymann, Valerie Beral, Matthew D. Snape, Helen Rees, Alba-Maria Ropero, Ran D. Balicer, Jakob P. Cramer, et al., "SARS-CoV-2 Variants and Vaccines," *New England Journal of Medicine* Special Report 385 (July 8, 2021):179–186; <https://doi.org/10.1056/NEJMSr2105280> (accessed July 9, 2021).
22. Brian J. Boyarsky, William A. Werbel, Robin K. Avery, et al., "Antibody Response to 2-Dose SARS-CoV-2 mRNA Vaccine Series in Solid Organ Transplant Recipients," *Journal of the American Medical Association* 325, no. 21 (May 5, 2021):2204–2206; <https://doi.org/10.1001/jama.2021.7489> (accessed July 9, 2021).
23. US Centers for Disease Control and Prevention, "When You've Been Fully Vaccinated: How to Protect Yourself and Others," updated June 17, 2021; <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/fully-vaccinated.html> (accessed July 9, 2021).
24. Matan Levine-Tiefenbrun, Idan Yelin, Rachel Katz, Esma Herzel, Ziv Golan, Licita Schreiber, Tamar Wolf, Varda Nadler, Amir Ben-Tov, Jacob Kuint, Sivan Gazit, Tal Patalon, Gabriel Chodick, and Roy Kishony, "Initial report of decreased SARS-CoV-2 viral load after inoculation with the BNT162b2 vaccine," *Nature Medicine* 27, no. 5 (March 29, 2021): 790–792; <https://doi.org/10.1038/s41591-021-01316-7> (accessed July 9, 2021).
25. Ibid.
26. Ross J. Harris, Jennifer A. Hall, Asad Zaidi, Nick J. Andrews, J. Kevin Dunbar, and Gavin Dabrera, "Effect of Vaccination on Household Transmission of SARS-CoV-2 in England," *New England Journal of Medicine*, published online June 23, 2021; <https://doi.org/10.1056/NEJMc2107717> (accessed July 9, 2021).
27. Yuhan Xing, Gary W. K. Wong, Ni Wei, Xiaowen Hu, and Quancheng Xing, "Rapid Response to an Outbreak in Qingdao, China," *New England Journal of Medicine* 383, no. 23 (November 18, 2020): e129; <https://doi.org/10.1056/NEJMc2032361> (accessed July 9, 2021).
28. Micheal J. Mina, Roy Parker, and Daniel B. Larremore, "Rethinking Covid-19 Test Sensitivity—A Strategy for Containment," *New England Journal of Medicine* 383, no. 22 (September 30, 2020): e120; <https://doi.org/10.1056/NEJMp2025631> (accessed July 9, 2021).
29. World Health Organization, "COVID-19 Virtual Press conference transcript—25 June 2021"; <https://www.who.int/publications/m/item/covid-19-virtual-press-conference-transcript-25-june-2021> (accessed July 9, 2021).
30. US Centers for Disease Control and Prevention, "SARS-CoV-2 Variant Classifications and Definitions," Updated June 29, 2021; <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html> (accessed July 9, 2021).
31. <https://apps.who.int/flumart/Default?ReportNo=7> (accessed July 9, 2021).
32. US Centers for Disease Control and Prevention, "Decreased Influenza Activity During the COVID-19 Pandemic—United States, Australia, Chile, and South Africa, 2020," *Morbidity and Mortality Report* 69, no. 37 (September 18, 2020):1305–1309; <https://www.cdc.gov/mmwr/volumes/69/wr/mm6937a6.htm> (accessed July 9, 2021).
33. Ali A. Rabaan, Shamsah H. Al-Ahmed, Maysaa K. Al-Malkey, Roua Alsubki, Sayeh Ezzikouri, Fadel Hassan Al-Hababi, et al., "Airborne transmission of SARS-CoV-2 is the dominant route of transmission: droplets and aerosols," *Le Infezioni in Medicina* 29, no. 1 (March 1, 2021): 10–19.
34. Sima Asadi, Nicole Bouvier, Anthony S. Wexler, and William D. Ristenpart, "The coronavirus pandemic and aerosols: Does COVID-19 transmit via expiratory particles?" *Aerosol Science and Technology* 54, no. 6 (June 2, 2020): 635–638; <https://doi.org/10.1080/02786826.2020.1749229> (accessed July 9, 2021).
35. Leonardo Setti, Fabrizio Passarini, Gianluigi De Gennaro, Perluigi Barbieri, Maria G. Perrone, Massimo Borelli, et al., "Airborne Transmission Route of COVID-19: Why 2 Meters/6 Feet of Inter-Personal Distance Could Not Be Enough," *International Journal of Environmental Research*

- and *Public Health* 17, no. 8 (April 23, 2020): 2932; <https://doi.org/10.3390/ijerph17082932> (accessed July 9, 2021).
36. Asadi, Bouvier, Wexler, and Ristenpart, 635–638.
 37. Ibid.
 38. Graham R. Johnson and Lidia Morawska, “The mechanism of breath aerosol formation,” *Journal of Aerosol Medicine and Pulmonary Drug Delivery* 22, no. 3 (September 2009): 229–237; <https://doi.org/10.1089/jamp.2008.0720> (accessed July 9, 2021).
 39. Asadi, Bouvier, Wexler, and Ristenpart, 635–638.
 40. Sima Asadi, Anthony S. Wexler, Christopher D. Cappa, Santiago Barreda, Nicole M. Bouvier, and William D. Ristenpart, “Aerosol Emission and Superemission during Human Speech Increase with Voice Loudness,” *Nature Scientific Reports* 9, no. 1 (February 20, 2019): 2348; <https://doi.org/10.1038/s41598-019-38808-z> (accessed July 9, 2021).
 41. Dirk Mürbe, Martin Kriegel, Julia Lange, Lukas Schumann, Anne Hartmann, and Mario Fleischer, “Aerosol emission of adolescents’ voices during speaking, singing and shouting,” *PLoS One* 16, no. 2 (February 10, 2021): e0246819; <https://doi.org/10.1371/journal.pone.0246819> (accessed July 9, 2021).
 42. Rabaan, Al-Ahmed, Al-Malkey, Alsubki, Ezzikouri, Al-Hababi, et al., 10–19.
 43. Wehzhao Chen, Nan Zhang, Jianjian Wei, Hui-Ling Yen, and Yuguo Li, “Short-range airborne route dominates exposure of respiratory infection during close contact,” *Building and Environment* 176 (June 2020): 106859; <https://doi.org/10.1016/j.buildenv.2020.106859> (accessed July 9, 2021).
 44. Ibid.
 45. Setti, Passarini, De Gennaro, Barbieri, Perrone, Borelli M, et al., 2932.
 46. Ibid.
 47. Ibid.
 48. Ibid.
 49. Joanna Zimmer-Nowicka and Hendryka Januszewska-Stańczyk, “Incidence and predisposing factors of common upper respiratory tract infections in vocal students during their professional training,” *Journal of Voice* 25, no. 4 (July 2011): 505–510; <https://doi.org/10.1016/j.jvoice.2010.01.015> (accessed July 9, 2021).
 50. Asadi, Wexler, Cappa, Barreda, Bouvier, and Ristenpart, 2348; Tehya Stockman, Shengwei Zhu, Abhishek Kumar, Lingzhe Wang, Sameer Patel, James Weaver, Mark Spede, Don Milton, Jean Hertzberg, Darin Toohey, Marina Vance, Jelena Srebric, and Shelly L. Miller, “Measurements and simulations of aerosol released while singing and playing wind instruments” [preprint, released April 15, 2021]; <https://scholar.colorado.edu/downloads/9s161736t> (accessed July 9, 2021); Mürbe, Kriegel, Lange, Schumann, Hartmann, and Fleischer, e0246819; Dirk Mürbe, Martin Kriegel, Julia Lange, Hansjorg Rotheudt, and Mario Fleischer, “Aerosol emission is increased in professional singing,” *Nature Scientific Reports* (in review); https://depositonce.tu-berlin.de/bitstream/11303/11491/5/muerbe_etal_2020_aerosols-singing.pdf (accessed July 9, 2021); John Volkens, Kristen Fedak, Dan Goble, Nick Good, Amy Kiesling, Christian L’Orange, Emily Morton, Rebecca Phillips, and Ky Tanner, “Update: bioaerosol emissions and exposures in the performing arts: a scientific roadmap for a safer return from COVID-19,” web presentation released May 20, 2021; https://smt.d.colostate.edu/wp-content/uploads/sites/34/2021/06/AerosolStudyPreliminaryResulits_05.21.2021.pdf (accessed July 10, 2021).
 51. National Association of Schools of Music, *NASM Handbook 2021* (Reston, VA: NASM, 2021); 67.
 52. American Society of Heating, Refrigerating and Air-Conditioning Engineers, *ANSI/ASHRAE Standard 62.2–2013: Ventilation and Acceptable Indoor Air Quality in Low-Rise Residential Buildings* (Atlanta: American Society of Heating, Refrigerating and Air-Conditioning Engineers, 2013), 19.
 53. Joseph G. Allen and Andrew M. Ibrahim, “Indoor Air Changes and Potential Implications for SARS-CoV-2 Transmission,” *Journal of the American Medical Association* 325, no. 20 (May 25, 2021): 2112–2113; <https://doi.org/10.1001/jama.2021.5053> (accessed July 10, 2021).
 54. Ibid.
 55. Tom Lipinski, Darem Ahmad, Nicholas Serey, and Hussam Jouhara, “Review of ventilation strategies to reduce the risk of disease transmission in high occupancy buildings,” *International Journal of Thermofluids* 7–8 (September 13, 2020): 100045; <https://doi.org/10.1016/j.ijft.2020.100045> (accessed July 10, 2021); US Centers for Disease Control and Prevention, “Ventilation in buildings (updated June 2, 2021);” <https://www.cdc.gov/coronavirus/2019-ncov/community/ventilation.html> (accessed July 10, 2021); American Society of Heating, Refrigerating and Air-Conditioning Engineers, “ASHRAE Position Document on Infectious Aerosols, approved by the ASHRAE Board of Directors April 14, 2020”; https://www.ashrae.org/file%20library/about/position%20documents/pd_infectiousaerosols_2020.pdf (accessed July 10, 2021).
 56. American Society of Heating, Refrigerating and Air-Conditioning Engineers, *ANSI/ASHRAE Standard 62.1–2019: Ventilation for Acceptable Indoor Air Quality* (Atlanta: ASHRAE; 2019), 10.
 57. American Society of Heating, Refrigerating and Air-Conditioning Engineers, “In-room air cleaner guidance for

- reducing COVID19 in air in your space/room” (accessed July 10, 2021).
58. US Environmental Protection Agency, “Indoor Air Quality: What is a HEPA filter?”; <https://www.epa.gov/indoor-air-quality-iaq/what-hepa-filter-1> (accessed July 10, 2021).
59. US Centers for Disease Control and Prevention, “Ventilation FAQs: What is a HEPA filter and why use a portable HEPA air cleaner”; <https://www.cdc.gov/coronavirus/2019-ncov/community/ventilation.html#refphf> (accessed July 10, 2021).
60. Zhe Peng and Jose L. Jimenez, “Exhaled CO₂ as a COVID-19 Infection Risk Proxy for Different Indoor Environments and Activities,” *Environmental Science and Technology Letters* 8, no. 5 (April 5, 2021): 392–397; <https://doi.org/10.1021/acs.estlett.1c00183> (accessed July 10, 2021); Lia Chatzidiakou, Dejan Mumovic, and Alex Summerfield, “Is CO₂ a good proxy for indoor air quality in classrooms? Part 1: The interrelationships between thermal conditions, CO₂ levels, ventilation rates and selected indoor pollutants,” *Building Services Engineering Research and Technology* 36, no. 2 (January 9, 2015): 129–161; <https://doi.org/10.1177/0143624414566244> (accessed July 10, 2021); Nisha Patel, Ciara Docherty, James R. Allison, Graham Walton, Ben Cole, Nicholas Jakubovics, Justin Durham, and Richard Holliday, “Letter to the Editor: Ventilation Is Key, Open a Window,” *OSF Preprints* (August 17, 2020); <https://doi.org/10.31219/osf.io/7rczy> (accessed July 10, 2021); Federation of European Heating, Ventilation and Air Conditioning Associations, “COVID-19 ventilation and building services guidance for school personnel”; https://www.rehva.eu/fileadmin/user_upload/REHVA_COVID-19_guidance_document_School_guidance_25112020.pdf (accessed July 10, 2021).
61. Angela Eykelbosh, “Can CO₂ sensors be used to assess COVID-19 transmission risk?” National Collaborating Centre for Environmental Health [blog, posted January 15, 2021]; <https://ncceh.ca/content/blog/can-co2-sensors-be-used-assess-covid-19-transmission-risk> (accessed July 10, 2021).
62. US Centers for Disease Control and Prevention, “Ventilation in Buildings; Ventilation FAQs: Can carbon dioxide (CO₂) monitors be used to indicate when there is good ventilation?” (updated June 2, 2021); <https://www.cdc.gov/coronavirus/2019-ncov/community/ventilation.html> (accessed July 10, 2021); Manfred Nusseck, Bernhard Richter, Ludwig Holtmeier, Dominik Skala, and Claudia Spahn, “CO₂ measurements in instrumental and vocal closed room settings as a risk reducing measure for a Coronavirus infection,” *medRxiv preprint* (October 27, 2020); <https://doi.org/10.1101/2020.10.26.20218354> (accessed July 10, 2021).
63. Ibid.
64. Stockman, Zhu, Kumar, et. al.
65. Ibid.; Nusseck, Richter, Holtmeier, Skala, and Spahn.
66. US Centers for Disease Control and Prevention Public Health Professionals’ Gateway, Public Health Law, Publications and Resources, “Health Insurance Portability and Accountability Act of 1996 (HIPAA); <https://www.cdc.gov/phlp/publications/topic/hipaa.html> (accessed July 8, 2021).
67. US Department of Education, “Family Educational Rights and Privacy Act (FERPA); <https://www2.ed.gov/policy/gen/guid/fpco/ferpa/index.html> (accessed July 8, 2021).
68. US Department of Education Student Privacy Policy Office “FERPA and Coronavirus Disease 2019 (COVID-19) Frequently Asked Questions (FAQs),” March, 2020; <https://studentprivacy.ed.gov/resources/ferpa-and-coronavirus-disease-2019-covid-19> (accessed July 8, 2021).
69. “FERPA and COVID-19 FAQs,” March, 2020, downloadable document pages 3–4; https://studentprivacy.ed.gov/sites/default/files/resource_document/file/FERPA%20and%20Coronavirus%20Frequently%20Asked%20Questions.pdf (accessed July 8, 2021).
70. Tom Bartlett, “At These Colleges, Professors Are Being Told Not to Ask Students if They’re Vaccinated,” *Chronicle of Higher Education*, June 2, 2021; <https://www.chronicle.com/article/at-these-colleges-professors-are-being-told-not-to-ask-students-if-theyre-vaccinated> (accessed July 8, 2021); Fernanda Zamudio-Suarez, “Weekly Briefing: Can You Talk About Vaccination Status to Students On Your Campus?” *Chronicle of Higher Education*, June 5, 2021; <https://www.chronicle.com/newsletter/weekly-briefing/2021-06-05> (accessed July 8, 2021).
71. National Association of Teachers of Singing “Joint Statement on SARS-CoV-2 Vaccinations and Best Health Practices,” June 23, 2021; https://www.nats.org/cgi/page.cgi/_article.html/What_s_New/Joint_Statement_on_SARS-CoV-2_Vaccinations_and_Best_Health_Practices (accessed July 8, 2021).
72. Jianyin Qiu, Bin Shen, Min Zhao, Zhen Wang, Bin Xie, and Yifeng Xu, “A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations,” *General Psychiatry* 33: e100213 (March 6, 2020); <http://dx.doi.org/10.1136/gpsych-2020-100213> (accessed July 10, 2021).
73. Maurizio Bonati, Rita Campi, Michele Zanetti, Massimo Cartabia, Francesca Scarpellini, Antonio Clavenna, and Giulia Segre, “Psychological distress among Italians during the 2019 coronavirus disease (COVID-19) quarantine,” *BMC Psychiatry* 21, no. 20 (January 8, 2021): 1–13; <https://doi.org/10.1186/s12888-020-03027-8> (accessed

- July 10, 2021); Anna Costantini and Eva Mazzotti, "Italian validation of CoViD-19 Peritraumatic Distress Index and preliminary data in a sample of general population," *Rivista di psichiatria* 55, no. 3 (May-June 2020): 145–151; <https://www.rivistadipsichiatria.it/archivio/3382/articoli/33570/> (accessed July 10, 2021).
74. Olga Megalakaki, Cyrille Kossigan Kokou-Kpolou, Justine Vaudé, Sunyoung Park, Steven Kator Iorfa, Jude Mary Cénat, and Daniel Derivois, "Does peritraumatic distress predict PTSD, depression and anxiety symptoms during and after COVID-19 lockdown in France? A prospective longitudinal study," *Journal of Psychiatric Research* 137 (May 2021): 81–88; <https://doi.org/10.1016/j.jpsychires.2021.02.035> (accessed July 10 2021).
 75. Roy Rillera Marzo, Emilio Quilatan Villanueva III, Erwin Martinez Faller, and Aries Moralidad Baldonado, "Factors associated with psychological distress among Filipinos during coronavirus disease-19 pandemic crisis," *Open Access Macedonian Journal of Medical Sciences* 8, no. T1 (October 28, 2020): 309–313; <https://doi.org/10.3889/oamjms.2020.5146> (accessed July 10, 2021).
 76. Asghar Afshar Jahanshahi, Maryam Mokhtari Dinani, Abbas Nazarian Madavani, Jizhen Li, and Stephen X. Zhang, "The distress of Iranian adults during the Covid-19 pandemic—More distressed than the Chinese and with different predictors," *Brain, Behavior, and Immunity* 87 (April 29, 2020): 124; <https://doi.org/10.1016/j.bbi.2020.04.081> (accessed July 10, 2021).
 77. Stephen X. Zhang, Yifei Wang, Asghar Afshar Jahanshahi, Jianfeng Jia, and Valentina Gomes Haensel Schmitt, "First study on mental distress in Brazil during the COVID-19 crisis" (January 17, 2021); <https://doi.org/10.1101/2020.04.18.20070896> (accessed July 10, 2021).
 78. Émilie Thomas, Daniel Saumier, and Alain Brunet, "Peritraumatic distress and the course of posttraumatic stress disorder symptoms: A meta-analysis," *The Canadian Journal of Psychiatry* 57, no. 2 (February 1, 2012): 122–129; <https://doi.org/10.1177/070674371205700209> (accessed July 10, 2021); Mary C. Vance, Benjamin Kovachy, Melissa Dong, and Eric Bui, "Peritraumatic distress: A review and synthesis of 15 years of research," *Journal of Clinical Psychology* 74, no. 9 (March 15, 2018): 1457–1484; <https://doi.org/10.1002/jclp.22612> (accessed July 10, 2021).
 79. Costantini and Mazzotti; Jahanshahi, Dinani, Madavani, Li, and Zhang; Shuyan Liu and Andreas Heinz, "Cross-cultural validity of psychological distress measurement during the coronavirus pandemic," *Pharmacopsychiatry* 53, no. 05 (June 24, 2020): 237–238; <https://pubmed.ncbi.nlm.nih.gov/32583390/> (accessed July 10, 2021).
 80. Cheng Cheng, Huang Jun, and Liang Baoyong, "Psychological health diathesis assessment system: a nationwide survey of resilient trait scale for Chinese adults," *Studies of Psychology and Behavior* 12, no. 6 (2014): 735; <http://psybeh.tjnu.edu.cn/EN/abstract/abstract898.shtml>.
 81. Duygu Kaba and Burcu Akin Sari, "Acute stress disorder with panic episodes induced by exposure to COVID-19 outbreak news in a child," *The Journal of Psychiatry and Neurological Sciences* 33 (April 13, 2020): 221–222; <http://dSPACE.baskent.edu.tr/bitstream/handle/11727/5808/1594121911-en.pdf?sequence=1&isAllowed=y> (accessed July 10, 2021).
 82. Costantini and Mazzotti; Qiu, Shen, Zhao, Wang, Xie, and Xu; Alberto Abad, Juliana Almeida da Silva, Lucas Emmanuel Pedro de Paiva Teixeira, Mayra Antonelli-Ponti, Sandra Bastos, Cláudia Helena Cerqueira Mármora, Luis Antonio Monteiro Campos, Scheila Paiva, Renato Leonardo de Freitas, and José Aparecido da Silva, "Evaluation of fear and peritraumatic distress during COVID-19 pandemic in Brazil," *Advances in Infectious Diseases* 10, no. 03 (September 14, 2020): 184; https://www.scirp.org/html/19-1950501_102814.htm (accessed July 10, 2021).
 83. Jitender Sareen, Julie Erickson, Maria I. Medved, Gordon JG Asmundson, Murray W. Enns, Murray Stein, William Leslie, Malcolm Doupe, Sarvesh Logsetty, "Risk factors for post-injury mental health problems," *Depression and Anxiety* 30, no. 4 (February 13, 2013): 321–327; <https://doi.org/10.1002/da.22077> (accessed July 10, 2021); Yong-Chun Bahk, Kiho Park, N. Kim, J. Lee, S. Cho, J. Jang, D. Jung, E. Chang, and K. Choi, "Psychological impact of COVID-19 in South Korea: a preliminary study," *Korean Journal of Clinical Psychology* 39, no. 4 (November 2020): 355–367; <https://doi.org/10.15842/kjcp.2020.39.4.008> (accessed July 10, 2021).
 84. Qiu, Shen, Zhao, Wang, Xie, and Xu; Zhang, Wang, Jahanshahi, Jia, and Haensel Schmitt.
 85. Cindy H. Liu, Emily Zhang, Ga Tin Fifi Wong, and Sunah Hyun, "Factors associated with depression, anxiety, and PTSD symptomatology during the COVID-19 pandemic: Clinical implications for US young adult mental health," *Psychiatry Research* 290 (August 2020): 113172; <https://doi.org/10.1016/j.psychres.2020.113172> (accessed July 10, 2021).
 86. Ibid.
 87. Wanjie Tang, Tao Hu, Baodi Hu, Chunhan Jin, Gang Wang, Chao Xie, Sen Chen, and Jiuping Xu, "Prevalence and correlates of PTSD and depressive symptoms one month after the outbreak of the COVID-19 epidemic in a sample of home-quarantined Chinese university students," *Journal of Affective Disorders* 274 (September 2020): 1–7; <https://doi.org/10.1016/j.jad.2020.05.009> (accessed July 10, 2021).

88. Abdelrahim Alqudah, Ahmad Al-Smadi, Muna Oqal, Esam Y. Qnais, Mohammed Wedyan, Majd Abu Gneam, Roaa Alnajjar, Manar Alajarmeh, Elaf Yousef, and Omar Gam-moh, "About anxiety levels and anti-anxiety drugs among quarantined undergraduate Jordanian students during COVID-19 pandemic," *International Journal of Clinical Practice* (April 21, 2021): e14249; <https://doi.org/10.1111/ijcp.14249> (accessed July 10, 2021).
89. Danny Horesh and Adam D. Brown, "Traumatic stress in the age of COVID-19: A call to close critical gaps and adapt to new realities," *Psychological Trauma: Theory, Research, Practice, and Policy* 12, no. 4 (March 25, 2020): 331; <https://doi.org/10.1037/tra0000592> (accessed July 10, 2021).
90. Ibid.
91. Jiaqi Xiong, Orly Lipsitz, Flora Nasri, Leanna M.W. Lui, Hartej Gill, Lee Phan, David Chen-Li, et al., "Impact of COVID-19 pandemic on mental health in the general population: A systematic review," *Journal of Affective Disorders* (December 1, 2020): 55–64; <https://doi.org/10.1016/j.jad.2020.08.001> (accessed July 10, 2021).
92. Elisa Monti, David C. Kidd, Linda M. Carroll, and Emanuele Castano, "What's in a singer's voice: the effect of attachment, emotions and trauma," *Logopedics Phoniatics Vocology* 42, no. 2 (April 14, 2016): 62–72; <https://doi.org/10.3109/14015439.2016.1166394> (accessed July 10, 2021).
93. Sharon Stohrer, *The Singer's Companion* (New York: Routledge Taylor and Francis Group, 2006), 103.
94. George S. Everly, Jr. and Jeffrey M. Lating, *A Clinical Guide to the Treatment of the Human Stress Response* (New York: Springer, 2019), 40–43.
95. Deborah Caputo Rosen, Reinhardt J. Heuer, David A. Sasso, and Robert T. Sataloff, "Psychologic Aspects of Voice Disorders," in Robert T. Sataloff, ed., *Diagnosis and Treatment of Voice Disorders* (San Diego: Plural Publishing, 2014), 599–600.
96. Robert Thayer Sataloff, "Stress, anxiety and psychogenic dysphonia," in Robert T. Sataloff, ed., *Professional Voice: The Science and Art of Clinical Care* (New York: Raven Press, 1991), 195–200.
97. Nora Siupsinskiene, Artūras Razbadauskas, and Laimis Dubosas, "Psychological distress in patients with benign voice disorders," *Folia Phoniatica et Logopædica* 63, no. 6 (March 24, 2011): 281–288; <https://doi.org/10.1159/000324641> (accessed July 10, 2021).
98. Judith Alyce Green and Robert Shellenberger, *The Dynamics of Health and Wellness: A Biopsychosocial Approach* (Orlando: Houghton Mifflin Harcourt School, 1991), 61–64, 92, 98, 101–136; Alan G. Stroudemire, *Psychological Factors Affecting Medical Conditions* (Washington, DC: American Psychiatric Press, 1995), 187–192.
99. Anna Vittoria Mattioli, Matteo Ballerini Puviani, Milena Nasi, and Alberto Farinetti, "COVID-19 pandemic: the effects of quarantine on cardiovascular risk," *European Journal of Clinical Nutrition* 74, no. 6 (May 5, 2020): 852–855; <https://doi.org/10.1038/s41430-020-0646-z> (accessed July 10, 2021).
100. Miriam van Mersbergen, "Research Approaches to the Intersection of Voice Disorders and Psychology," in *Psychology of Voice Disorders* (San Diego: Plural Publishing, 2020), 53–54.
101. Amanda I. Gillespie, Leah B. Helou, and Aaron Ziegler, "Crossroads of the respiratory and phonatory systems: where to go from here?" [Conference presentation] 43rd Annual Symposium on the Care of the Professional Voice (Philadelphia PA, May 30, 2014).
102. Peter Butcher, Annie Elias, and Lesley Cavalli, *Understanding and Treating Psychogenic Voice Disorder: A CBT Framework* (Hoboken: John Wiley & Sons, 2007).
103. Avi Besser, Sari Lotem, and Virgil Zeigler-Hill, "Psychological stress and vocal symptoms among university professors in Israel: implications of the shift to online synchronous teaching during the COVID-19 pandemic," *Journal of Voice* (in press; available July 5, 2020); <https://doi.org/10.1016/j.jvoice.2020.05.028> (accessed July 10, 2021).
104. Monti, Kidd, Carroll, and Castano.
105. Bessel Van der Kolk, *The Body Keeps the Score: Brain, Mind, and Body in the Healing of Trauma* (New York: Penguin Books, 2014): 108.
106. Ibid.
107. Stephen W. Porges, "The polyvagal theory: phylogenetic substrates of a social nervous system," *International Journal of Psychophysiology* 42, no. 2 (October, 2001):144; [https://doi.org/10.1016/S0167-8760\(01\)00162-3](https://doi.org/10.1016/S0167-8760(01)00162-3) (accessed July 10, 2021).
108. Stephen W. Porges, "Polyvagal theory: A primer," in *Clinical Applications of the Polyvagal Theory: The Emergence of Polyvagal-informed Therapies* (Norton Series on Interpersonal Neurobiology) (New York: W.W. Norton & Company, 2018), 67.
109. Ibid., 61.
110. Dianna Kenny, *The Psychology of Music Performance Anxiety* (Oxford: Oxford University Press, 2011).
111. Lynn Holding, *The Musician's Mind: Teaching, Learning, and Performance in the Age of Brain Science* (Lanham, MD: Rowman & Littlefield, 2020): 207–258.

112. US Department of Health and Human Services, "Clinical Practice Guideline Number 5: Depression in Primary Care, vol 2: Treatment of Major Depression: AHCPR Publication 93-0550," (Rockville, MD: US Department of Health and Human Services Agency for Health Care Policy and Research, 1993).
113. Rosen, Heuer, Sasso, and Sataloff.
114. Robert A. Bell, Debora A. Paterniti, Rahman Azari, Paul R. Duberstein, Ronald M. Epstein, Aaron B. Rochlen, Megan Dwight Johnson, Sharon E. Orrange, Christina Slee, and Richard L. Kravitz, "Encouraging patients with depressive symptoms to seek care: a mixed methods approach to message development," *Patient Education and Counseling* 78, no. 2 (February, 2010): 198-205; <https://doi.org/10.1016/j.pec.2009.07.007> (accessed July 10, 2021); Georg Schomerus and Matthias C. Angermeyer, "Stigma and its impact on help-seeking for mental disorders: what do we know?" *Epidemiology and Psychiatric Sciences* 17, no. 1 (March 2008): 31-37; <https://doi.org/10.1017/S1121189X00002669> (accessed July 10, 2021); Stephen P. Hinshaw and Dante Cicchetti, "Stigma and mental disorder: Conceptions of illness, public attitudes, personal disclosure, and social policy," *Development and Psychopathology* 12, no. 4 (December 2000): 555-598; <https://doi.org/10.1017/S0954579400004028> (accessed July 10, 2021); Amelia Gulliver, Kathleen M. Griffiths and Helen Christensen, "Perceived barriers and facilitators to mental health help-seeking in young people: a systematic review," *BMC Psychiatry* 10, no. 113 (December 30, 2010): 1-9; <https://doi.org/10.1186/1471-244X-10-113> (accessed July 10, 2021).
115. Brandy R. Maynard, Anne Farina, Nathaniel A. Dell, and Michael S. Kelly, "Effects of trauma-informed approaches in schools: A systematic review," *Campbell Systematic Reviews* 15, no. 1-2 (2019): e1018; <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cl2.1018> (accessed July 10, 2021).
116. Kristen Doughty, "Increasing Trauma Informed Awareness in Higher Education" (EdD dissertation, Wilmington University, 2018), 39-40.
117. Grace Weber Johnson, "The Private Studio-Singing and Childhood Sexual Abuse: Conversation with Voice Teachers," *Journal of Singing* 65, no. 4 (March/April 2009): 471-480; Lauren McQuistin, "The Effects of Childhood Sexual Abuse on Singers," *Journal of Singing* 76, no. 4 (March/April 2020): 423-428; Jacqueline Robarts, "Music therapy with sexually abused children," *Clinical Child Psychology and Psychiatry* 11, no. 2 (April 1, 2006): 249-269; <https://doi.org/10.1177%2F1359104506061418> (accessed July 10, 2021); Catherine Y. Wan, Theodor Rüber, Anja Hohmann, and Gottfried Schlaug, "The therapeutic effects of singing in neurological disorders," *Music Perception* 27, no. 4 (April 1, 2010): 287-295; <https://doi.org/10.1525/mp.2010.27.4.287> (accessed July 10, 2021).
118. Jess Baldwin, "Incorporating Basic Trauma Awareness into the Voice Studio"; <https://singinginpopularmusics.com/2020/04/06/incorporating-basic-trauma-awareness-into-the-voice-lesson/> (accessed July 1, 2021).
119. I. M. Van Eekelen, H. P. A. Boshuizen, and Jan D. Vermunt, "Self-regulation in higher education teacher learning," *Higher Education* 50, no. 3 (October 2005): 447-471; <https://doi.org/10.1007/s10734-004-6362-0> (accessed July 10, 2021).
120. Laura Crocco and David Meyer, "Motor Learning and Teaching Singing: An Overview," *Journal of Singing* 77, no. 5 (May/June 2021): 693-702.
121. David Meyer and Lynn Holding, "Practical Science in the Studio: 'No-Tech' Strategies," *Journal of Singing* 77, no. 3 (January/February 2021): 359-367.
122. Lynn Holding, Thomas L. Carroll, John Nix, Michael M. Johns, Wendy D. LeBorgne, and David Meyer, "COVID-19 after effects: concerns for singers," *Journal of Voice* (in-press, August 6, 2020); <https://doi.org/10.1016/j.jvoice.2020.07.032> (accessed July 10, 2021).
123. Roberto Burioni and Eric J. Topol, "Has SARS-CoV-2 reached peak fitness?" *Nature Medicine* (June 21, 2021): 1; <https://doi.org/10.1038/s41591-021-01421-7> (accessed July 10, 2021).

A leading scholar and researcher of the singing voice, **Dr. David Meyer** is an active performer, teacher, clinician, and voice scientist. He serves as associate professor of voice and voice pedagogy at Shenandoah Conservatory, and is Director of the Janette Ogg Voice Research Center. He is also a member of the Scientific Advisory Board of the Voice Foundation and is Co-chairman of the Voice Science Advisory Committee of the National Association of Teachers of Singing. In 2010 he received the Van L. Lawrence Fellowship, a prestigious national award in recognition of his contributions to the field of teaching singing and the use of voice science. Dr. Meyer's students have won numerous awards and have sung in major venues worldwide. Please see www.davidmeyervoice.com for more information.

John Nix is Professor of Voice and Voice Pedagogy and chair of the voice area at the University of Texas at San Antonio. His mentors include Barbara Doscher (singing, pedagogy) and Ingo Titze (voice science). Current and former students have sung with the Santa Fe, Arizona, Chautauqua, St. Louis, Nevada, Omaha, and San Antonio opera companies, and two have served as NATS Intern Program Master Teachers. In addition to his active voice teaching studio, he performs research in voice pedagogy, literature, and acoustics, having produced 46 published articles and 8 book chapters; he also co-chairs the NATS Voice Science Advisory Committee, and

serves on the *Journal of Singing's* editorial board. Mr. Nix is editor and annotator of *From Studio to Stage: Repertoire for the Voice* (Scarecrow, 2002), vocal music editor for the *Oxford Handbook of Music Education* (Oxford University Press, 2012), and one of three general editors for the *Oxford Handbook of Singing* (Oxford University Press, 2019).

Lynn Holding is Coordinator of Vocology and Voice Pedagogy and a studio voice teacher at the USC Thornton School of Music. She is the author of *The Musician's Mind: Teaching, Learning & Performance in the Age of Brain Science*, the chapter "Brain" in *Your Voice: An Inside View* 3rd ed. by Scott McCoy, and has served as an associate editor of the *Journal of Singing*, where she created the "Mindful Voice" column in the *Journal of Singing*, authoring it from its debut in October 2009 to the final installment in October 2017. Holding's voice science honors include the 2005 *Van L. Lawrence Fellowship*, and election to chair the founding of the first non-profit vocology association PAVA, incorporated in 2014 as a 501(c)(6) non-profit association. Holding's stage credits include leading roles with Harrisburg Opera, Nashville Opera, and Ohio Light Opera. She has commissioned new works and refashioned traditional recitals into theatrical performance pieces presented throughout the United States, Australia, England, France, Germany, Italy, Spain, and Iceland. Holding studied voice at the University of Montana with Esther England, in Vienna with Otto Edelmann, and at Indiana University with Dale Moore, where she was the first singer to pursue the Artist Diploma. She earned her master's degree in voice pedagogy from Westminster Choir College of Rider University and completed the Summer Vocology Institute at the National Center for Voice and Speech. Please see www.lynnholding.com for more information.

Allen Henderson holds degrees from Carson Newman College (BM), The University of Tennessee (MM), and the College-Conservatory of Music at the University of Cincinnati (DMA), where his minor was in Arts Administration and he was winner of the prestigious Corbett-Treigle Opera Competition. Prior to his appointment as executive director of NATS in 2008, he served as district and regional governor and was elected national secretary/treasurer from 2006–2008. He also served as interim executive director from 2007–2008. Henderson was a participant in the 1993 NATS Intern Program and later hosted the program in 1998. He was chair of the coordinating committee for the 2008 NATS 50th National Conference in Nashville, Tenn. Henderson has held teaching positions at Oklahoma Baptist University, Austin Peay State University, and Georgia Southern University. In these positions he has taught voice, foreign language diction, opera, choral techniques, choral literature, song literature, and directed choirs. He also served as music department chair at Austin Peay and Georgia Southern. As baritone soloist, Henderson has appeared in concert, opera, and oratorio across the country. A district winner and regional finalist in the Metropolitan Opera auditions, Henderson was winner of the 1995 National Federation of Music Clubs Artist Awards. He also was a regional finalist in the NATSAA competition.

Dr. Thomas L. Carroll is a surgeon at Brigham and Women's Hospital (BWH) specializing in laryngology and Assistant Professor of Otolaryngology at Harvard Medical School. Since 2014 he has served as the director of the BWH Voice Program, which provides diagnoses and innovative therapies for patients with voice, swallowing, and airway disorders. Dr. Carroll's interest in voice began as a boy chorister at St. Thomas Choir School in New York and continued through a music degree at Oberlin College. He received his MD from Wright State University School of Medicine and completed his residency in otolaryngology at the University of Colorado-Denver and Health Sciences Center. Dr. Carroll completed a fellowship in laryngology and care of the professional voice at the University of Pittsburgh Medical Center Voice Center. Clinical interests include care of the professional voice, early glottic cancer with an emphasis on voice preservation, laryngopharyngeal reflux, and related disorders such as chronic cough, vocal cord paralysis/paresis, and office-based laryngeal surgery, including vocal fold augmentation for both diagnostic and therapeutic purposes and photoangiolytic (KTP) laser therapy.

Jeremy Samuel Faust, MD, MS, MA (@jeremyfaust) is an attending physician in the Brigham and Women's Hospital Department of Emergency Medicine in the Health Policy Division and is an Instructor at Harvard Medical School. He is medical editor-in-chief of *ACEP Now* and an associate editor of News & Perspectives for *The Annals of Emergency Medicine*. His writing has appeared in *The New York Times*, *Slate*, *The New York Daily News*, *Mother Jones*, *Undark*, and peer reviewed journals including *JAMA*, *Lancet Oncology*, *The Annals of Emergency Medicine*, and others. He has spoken internationally at the Royal College of Emergency Medicine, Social Media and Critical Care (Australia, Germany, Ireland), and domestically at the Counsel of Residency Directors of Emergency Medicine (CORD); Harvard Medical School's Writing, Publishing, and Social Media for Healthcare Professionals; Resuscitation; and elsewhere. He and Lauren Westafer, DO, MPH, are the cohosts of the award winning FOAMcast, a popular emergency medicine podcast and inaugural recipient of EMRA's FOAMer of the Year award.

Dr. Christine Petersen's scholarly work has focused on the recognition and prevention of zoonotic diseases, primarily the epidemiology and immunobiology of vector-borne and parasitic diseases, and now, SARS-CoV-2. Dr. Petersen is the scientific program chair for the American Society of Tropical Medicine and Hygiene. Her collaborative group works in Brazil, India and Ethiopia via NIH/Fogarty International Center and NIAID-funded studies focused on understanding transmission and host immune susceptibility for zoonoses. Dr. Petersen's published and patented work demonstrates the ability to target reservoir species for immunologic and parasitologic control of infections to promote disease elimination. As director of the Center for Emerging Infectious Diseases, Dr. Petersen coordinates One Health activities focused on understanding, detecting and preventing emerging zoonotic disease globally. Dr. Petersen is a Professor at University of Iowa, College of Public Health, Department of Epidemiology.